arXiv:1607.00132v1 [math.MG] 1 Jul 2016

REDUCED SPHERICAL CONVEX BODIES

MAREK LASSAK and MICHAŁ MUSIELAK

Institute of Mathematics and Physics, University of Science and Technology, Kaliskiego 7, Bydgoszcz 85-789, Poland
e-mails: marek.lassak@utp.edu.pl, michal.musielak@gmail.com

Abstract

The aim of this paper is to present some properties of reduced spherical convex bodies on the two-dimensional sphere S^2. The intersection of two different non-opposite hemispheres is called a lune. By its thickness we mean the distance of the centers of the two semicircles bounding it. The thickness $\Delta(C)$ of C is the minimum thickness of a lune containing C. We say that a spherical convex body R is reduced if $\Delta(Z) < \Delta(R)$ for every spherical convex body $Z \subset R$ different from R. Our main theorem permits to describe the shape of reduced bodies of thickness below $\frac{\pi}{2}$. It implies a number of corollaries. In particular, we estimate the diameter of reduced spherical bodies in terms of their thickness. Reduced bodies of thickness at least $\frac{\pi}{2}$ have constant width. Spherical convex bodies of constant width below $\frac{\pi}{2}$ are strictly convex.

1 Introduction

Let S^d be the unit sphere of the $(d+1)$-dimensional Euclidean space E^{d+1}. By a *great circle* of S^d we mean the intersection of S^d with any two-dimensional subspace of E^{d+1}. The common part of S^d with any hyper-subspace of E^{d+1} is called a $(d-1)$-*dimensional great sphere* of S^d. In particular, for S^2 it is nothing else but a great circle. By a pair of *antipodes* of S^d we mean any pair of points which are obtained as the intersection of S^d with a one-dimensional subspace of E^{d+1}. Observe that if two different points are not antipodes, there is exactly one great circle containing them. If different points $a, b \in S^d$ are not antipodes, by the *spherical arc*, or shortly *arc*, ab connecting them we understand the shorter part of the great circle containing a and b. By the *spherical distance* $|ab|$, or shortly *distance*, of these points we mean the length of the arc connecting them. A subset of S^d is called *convex* if it does not contain any pair of antipodes of S^d and if together with every two points it contains the arc connecting them. By *a spherical convex body* we mean a closed convex set with non-empty interior. If in the boundary of a spherical convex body there is no arc, we say that the body is *strictly convex*. Convexity on S^d is considered in very many papers and monographs. For instance in [1], [2], [3] [4], [5], [7], [16], [17], [18].

The set of points of S^d in the distance at most ρ, where $\rho \in (0, \frac{\pi}{2})$, from a point $c \in S^d$ is called a *spherical ball*, or shorter *a ball*, of *radius* ρ and *center* c. Balls on S^2 are called *disks* and the boundary of a disk is called a *spherical circle*. Balls of radius $\frac{\pi}{2}$ are called *hemispheres*. The hemisphere with center m is denoted by $H(m)$. In other words, a *hemisphere* is the

Key words and phrases: spherical convex body, spherical geometry, hemisphere, lune, width, constant width, thickness, diameter, reduced body, extreme point

Mathematics Subject Classification: Primary 52A55; Secondary 97G60

intersection of S^d with any closed half-space of E^{d+1}. Two hemispheres whose centers are antipodes are called *opposite hemispheres*.

Let p be a boundary point of a convex body $C \subset S^d$. We say that a hemisphere H *supports* C *at* p provided $C \subset H$ and p belongs to the $(d-1)$-dimensional great sphere bounding H. If at p the body C is supported by exactly one hemisphere, we say that p *is a smooth point of* C. If all boundary points of C are smooth, then C is called to be *smooth*.

A *spherical* $(d-1)$*-dimensional ball of radius* $\rho \in \left(0, \frac{\pi}{2}\right]$ is the set of points of a $(d-1)$-dimensional great sphere of S^d in the distance at most ρ from a fixed point, called *the center* of this ball. The $(d-1)$-dimensional balls of radius $\frac{\pi}{2}$ are called $(d-1)$*-dimensional hemispheres*, and if $d = 2$ we call them *semicircles*.

If hemispheres G and H are different and not opposite, then $L = G \cap H$ is called a *lune*. The two $(d-1)$-dimensional hemispheres bounding L and contained in G and H, respectively, are denoted by G/H and H/G. The *thickness* $\Delta(L)$ of $L \subset S^d$ is defined as the distance of the centers of G/H and H/G. For $d = 2$ the $(d-1)$-dimensional hemispheres bounding L are called the *semicircles bounding* L. For $d = 2$, by the *corners* of L we understand the two points of the set $(G/H) \cap (H/G)$.

For every hemisphere K supporting a convex body $C \subset S^d$ we are finding hemispheres K^* supporting C such that the lunes $K \cap K^*$ are of the minimum thickness (by compactness arguments at least one such a hemisphere K^* exists). The thickness of the lune $K \cap K^*$ is called *the width of* C *determined by* K and it is denoted by $\mathrm{width}_K(C)$ (see [12]). The width of C determined by K changes continuously, as the position of K changes (see Theorem 2 of [12]). If for all hemispheres K supporting C the numbers $\mathrm{width}_K(C)$ are equal, we say that C is *of constant width* (see [12]). An application of spherical convex bodies of constant width is given in [8].

By the *thickness* $\Delta(C)$ of a convex body $C \subset S^d$ we understand the minimum width of C determined by K over all supporting hemispheres K of C (see [12]). The thickness of C is nothing else but the minimum thickness of a lune containing C (see [12]); this approach is sometimes more convenient.

After [12] we say that a spherical convex body $R \subset S^d$ is *reduced* if $\Delta(Z) < \Delta(R)$ for every convex body $Z \subset R$ different from R. This definition is analogical to the definition of a reduced body in Euclidean space E^d given in [9]. See also [6], [11] and the survey article [14]. For a larger context see Part 5.4 of [10]. Simple examples of reduced spherical convex bodies on S^d are spherical bodies of constant width and, in particular, the balls on S^d. Also each of the 2^d parts of a spherical ball on S^d dissected by d great pairwise orthogonal $(d-1)$-dimensional spheres through the center of a spherical ball B is a reduced spherical body. It is called $\frac{1}{2^d}$*-part of a ball* (see [12]). In particular, for $d = 2$, it is called a *quarter of a spherical disk*. There is a wide class of reduced odd-gons on S^2 (see [13]). In particular, spherical regular odd-gons of thickness at most $\frac{\pi}{2}$ are reduced. Figure 2 of [13] shows a non-regular reduced pentagon.

2 Supporting hemispheres of a spherical convex body

Let $C \subset S^2$ be a spherical convex body and let X, Y, Z be different supporting hemispheres of C. Denote by x, y, z the centers of X, Y, Z, respectively. We introduce the following three-argument relationship $\prec XYZ$. In the case when there is no common point of the three great circles bounding X, Y, Z and if x, y, z are in this order on the boundary of the spherical triangle xyz according to the positive orientation, then we write $\prec XYZ$. In the case when the three great circles bounding X, Y, Z have a common point c and if x, y, z are in this order on the boundary of the hemisphere centered at c, then we write $\prec XYZ$. If $\prec XYZ$, then we say that X, Y, Z *support* C *in this order*. We need the notion $\prec XYZ$ as a tool in the proof of our main Theorem 1, and also for Lemma 2 applied in its proof.

By the way, $\prec XYZ$ if and only if $\prec YZX$ and if and only if $\prec ZXY$.

The symbol $\preceq XYZ$ means that $\prec XYZ$ or $X = Y$ or $Y = Z$ or $Z = X$. If $\prec XYZ$ (respectively: $\preceq XYZ$), then we say that Y *supports* C *strictly between* (respectively: *between*) X *and* Z.

Assume that $\prec XYZ$, where X, Y, Z are supporting hemispheres of C. Then if we go on the boundary of C according to the positive orientation starting at any point of $C \cap \mathrm{bd}(X)$, we do not reach any point of $C \cap \mathrm{bd}(Z)$ before reaching all points of $C \cap \mathrm{bd}(Y)$.

Observe that for any boundary point p of a spherical convex body $C \subset S^2$ there are hemispheres H_r and H_l supporting C at p such that every hemisphere H supporting C at p satisfies $H_r \preceq H \preceq H_l$. We call H_r and H_l the *right* and, respectively, *the left supporting hemispheres of* C *at* p. Clearly, if p is a smooth point of C, then the hemispheres H_r and H_l coincide. The left and right supporting hemispheres of C are called *extreme supporting hemispheres of* C.

Lemma 1. *Let $C \subset S^2$ be a convex body and let $L = G \cap H$ be a lune of thickness $\Delta(C)$ containing C, where G and H are hemispheres. Denote by g the center of G/H and by h the center of H/G. We claim that at least one of the hemispheres G and H is a right supporting hemisphere of R (at g if this is G and at h if this is H), and at least one of these hemispheres is a left supporting hemisphere of C (at g if this is G and at h if this is H).*

The proof is analogous to the proof of Proposition 2.1 from [13].

Having in mind the forthcoming Lemma 2 and Theorem 1, following Part I of Theorem 1 of [12] recall that for every supporting hemisphere N of a spherical convex body $C \subset S^2$ of thickness below $\frac{\pi}{2}$ the hemisphere N^* is unique.

Lemma 2. *Assume that hemispheres N_1, N_2 and N_3 support a spherically convex body C of thickness below $\frac{\pi}{2}$ and let $\mathrm{width}_{N_i}(C) = \Delta(C)$ for $i = 1, 2, 3$. We have $\prec N_1N_2N_3$ if and only if $\prec N_1^*N_2^*N_3^*$.*

Proof. Since $\mathrm{width}_{N_i}(C) = \Delta(C)$, the thickness of the lune $N_i \cap N_i^*$ is $\Delta(C)$ for $i = 1, 2, 3$. Denote by a_i the center of the semicircle N_i/N_i^* and by b_i the center of the semicircle N_i^*/N_i for $i = 1, 2, 3$. By Claim 2 of [12] these six centers belong to the boundary of C.

Denote by o_i be the center of the hemisphere N_i and by o_i^* the center of the hemisphere N_i^* for $i = 1, 2, 3$. Since $\Delta(C) < \frac{\pi}{2}$, we see that o_i and o_i^* do not belong to C for $i = 1, 2, 3$. Thus a_i and b_i belong to the arc $o_io_i^*$ for $i = 1, 2, 3$.

Put $A = \{a_1, a_2, a_3, b_1, b_2, b_3\}$ and $O = \{o_1, o_2, o_3, o_1^*, o_2^*, o_3^*\}$. Observe that every point of A belongs to any hemisphere with the center in O. Thus the distance between any point of O and any point of A is at most $\frac{\pi}{2}$. Consequently, every hemisphere with the center in A contains O.

Denote by V the intersection of the six hemispheres with the centers in A. From the inclusion $C \subset V$ we conclude that V has non-empty interior. Moreover, the points of A are not contained in one great circle, which implies that V does not contain any pair of antipodes. Consequently, V is a spherical convex body. Observe that all points of O belong to the boundary of V.

Note that the length of $o_io_i^*$ is equal $\pi - \Delta(R)$ and is greater than $\frac{\pi}{2}$ for $i = 1, 2, 3$. Consider the lune $L_1 = H(a_1) \cap H(b_1)$. By $V \subset L_1$ we conclude that all points from O belong to L_1. Denote by c and c' corners of L_1 and consider two spherical triangles $T = o_1co_1^*$ and $T' = o_1c'o_1^*$. Clearly, $T \cup T' = L_1$. Note that $\mathrm{diam}(T) = \mathrm{diam}(T') = \pi - \Delta(R)$ and the only arc of length $\pi - \Delta(R)$ contained in T or contained in T' is $o_1o_1^*$. Therefore either $o_1 = o_2^*$ and $o_1^* = o_2$, or exactly one of points o_2 and o_2^* belongs to T and exactly one belongs to T'. Thus either $o_1o_1^* = o_2^*o_2$ or $o_1o_1^*$ intersects $o_2o_2^*$ and the point of the intersection belongs to the interior of V. In the same way we show that either two arcs from amongst $o_1o_1^*$, $o_2o_2^*$,

$o_3o_3^*$ coincide, or each pair of them intersects in the interior of V. In both cases we conclude that o_1, o_2, o_3 are on the boundary of V in this order according to the positive orientation if and only if o_1^*, o_2^*, o_3^* lay on the boundary V in this order according to the positive orientation. Thus by the convexity of V we obtain that $\prec N_1N_2N_3$ if and only if $\prec N_1^*N_2^*N_3^*$. □

3 Reduced bodies of thickness below $\frac{\pi}{2}$

Our main Theorem 1 gives a description of the shape of reduced bodies of thickness below $\frac{\pi}{2}$ on S^2. It is analogous to its version in E^2 presented in Theorem 3 of [11]. We are not able to present analogical proof as this in [11]. The first trouble is in the lack of the notion of parallelism on the sphere, which was used many times in the proof for E^2. The second trouble is that we are not able to give a spherical analog of the relationship $l_1 \prec l_2$ for a pair of different directions l_1, l_2 considered in [11] for E^2. It appears that a similar notion $G_1 \prec G_2$ for pairs of great circles, or a notion $H_1 \prec H_2$ for pairs of hemispheres (even supporting a spherical convex body C) does not make sense; the reason is that every two different great circles intersect at two points (so not at one point as two different non-parallel straight lines in the plane), and thus we are not able to state which of these great circles is "earlier". What helps, is the notion of the three argument relationship $\prec XYZ$ presented in Section 2. It permits to establish the order of any three different hemispheres X, Y, Z supporting a spherical convex body C, which appears to be sufficient for our needs in the proof of Theorem 1 and also the applied there Lemma 2.

Theorem 1. *Let $R \subset S^2$ be a reduced spherical body with $\Delta(R) < \frac{\pi}{2}$. Let M_1 and M_2 be supporting hemispheres of R such that* $\mathrm{width}_{M_1}(R) = \Delta(R) = \mathrm{width}_{M_2}(R)$ *and* $\mathrm{width}_M(R) > \Delta(R)$ *for every hemisphere M satisfying $\prec M_1MM_2$. Consider the lunes $L_1 = M_1 \cap M_1^*$ and $L_2 = M_2 \cap M_2^*$. We claim that the arcs a_1a_2 and b_1b_2 are in the boundary of R, where a_i is the center of M_i/M_i^* and b_i is the center of M_i^*/M_i for $i = 1, 2$. Moreover, $|a_1a_2| = |b_1b_2|$.*

Proof. By Claim 2 of [12] points a_1, b_1, a_2, b_2 belong to R and thus to the boundary of R. If we assume that the piece of the boundary of R from a_1 to a_2 (according to the positive orientation) is not an arc, then there is an extreme point e of R on this piece of the boundary different from a_1 and a_2. Then by Theorem 4 from [12] there exists a lune $L \supset R$ of thickness $\Delta(R)$ with e as the center of one of the two semicircles bounding L. Denote by M_L the hemisphere for which $e \in \mathrm{bd}(M_L)$, where $L = M_L \cap M_L^*$. Observe that L differs from L_1 and L_2. The reason is that if we assume $L = L_i$ for $i \in \{1, 2\}$, then both a_i and e are the centers of the same semicircle bounding the lune $L = L_i$, and thus $a_i = e$, which is impossible. Since L differs from L_1 and L_2, and since e belongs to the piece of the boundary from a_1 to a_2 being different from these points, we have $\prec M_1M_LM_2$. This, the description of M_L and $\Delta(L) = \Delta(R)$ contradict the assumption of our theorem that for every hemisphere M satisfying $\prec M_1MM_2$ we have $\mathrm{width}_M(R) > \Delta(R)$. Hence the considered piece of the boundary must be the arc a_1a_2.

Now we intend to show that the piece of the boundary of R from b_1 to b_2 (according to the positive orientation) is the arc b_1b_2. Assume the opposite. Then there exists an extreme point e' of R on this piece different from b_1 and b_2. Consequently, by Theorem 4 from [12] there exists a lune $L' \supset R$ of thickness $\Delta(R)$ with e' as the center of one of the two hemispheres bounding L'. Moreover, L' has the form $M_{L'} \cap (M_{L'})^*$ where $\prec M_1^*(M_{L'})^*M_2^*$.

By Lemma 2 we have $\prec M_1M_{L'}M_2$. This and $\mathrm{width}_{M_{L'}}(R) = \Delta(R)$ contradict the assumption of our theorem that $\mathrm{width}_M(R) > \Delta(R)$ for every hemisphere M fulfilling $\prec M_1MM_2$. Consequently the considered piece of the boundary of R is the arc b_1b_2.

Finally, show that $|a_1a_2| = |b_1b_2|$.



FIGURE 1. Illustration to the proof of Theorem 1.

Let x, y, z, w be the vertices of the spherical quadrangle $L_1 \cap L_2$ such that $a_1 \in wx, a_2 \in xy, b_1 \in yz$ and $b_2 \in zw$. Denote by H_a the hemisphere supporting R at every point of the arc a_1a_2 and by H_b the hemisphere supporting R at every point of the arc b_1b_2. Observe that H_a is the left supporting hemisphere of R at a_1 and the right supporting hemisphere of R at a_2. Analogously, H_b is the left supporting hemisphere of R at b_1 and the right supporting hemisphere of R at b_2. From Lemma 1 we know that at least one of the hemispheres M_1 and M_1^* is a left supporting hemisphere of R at a boundary point of R. Therefore $M_1 = H_a$ or $M_1^* = H_b$, and for the same reason we have $M_2 = H_a$ or $M_2^* = H_b$.

We cannot have simultaneously $M_1 = H_a$ and $M_2 = H_a$ because M_1 is different from M_2. For a similar reason it cannot be simultaneously $M_1^* = H_b$ and $M_2^* = H_b$. Hence either $M_1 = H_a$ and $M_2^* = H_b$, or $M_1^* = H_b$ and $M_2 = H_a$. As a consequence either $a_1 = x$ and $b_2 = z$, or $a_2 = x$ and $b_1 = z$.

Let for instance $a_1 = x$ and $b_2 = z$ (see Figure 1). By the assumptions of our theorem we have $|a_1b_1| = \Delta(R) = |a_2b_2|$. Moreover, the triangles $a_1a_2b_2$ and $b_2b_1a_1$ have the common side a_1b_2 and equal angles $\angle a_1a_2b_2 = \frac{\pi}{2} = \angle b_2b_1a_1$ (see Figure 1). Hence they are equal. Thus $|a_1a_2| = |b_1b_2|$. □

Theorem 1 allows us to describe the structure of the boundary of every reduced body on S^2. Namely, we conclude that the boundary consists of in a sense "opposite" arcs of equal length (some pairs of successive two such arcs may form longer arcs) and from some "opposite" pieces of spherical "curves of constant width". We obtain these "opposite" pieces of spherical curves of constant width always when $\mathrm{width}_M(R) = \Delta(R)$ for all M fulfilling $\preceq M_1MM_2$, where M_1 and M_2 are two fixed supporting semicircles of R.

Proposition 1. *Let $R \subset S^2$ be a reduced body with $\Delta(R) < \frac{\pi}{2}$. Assume that $\mathrm{width}_{M_1}(R) = \Delta(R) = \mathrm{width}_{M_2}(R)$, where M_1 and M_2 are two fixed supporting hemispheres of R. Denote by a_i the center of the semicircle M_i/M_i^*, and by b_i the center of the semicircle M_i^*/M_i for $i = 1, 2$. Assume that $a_1 = a_2$ and that $\preceq M_1MM_2$ for every M supporting R at this point. Then the shorter piece of the spherical circle with the center $a_1 = a_2$ and radius $\Delta(R)$ connecting b_1 and b_2 is in the boundary of R. Moreover, $\mathrm{width}_M(R) = \Delta(R)$ for all these M.*

Proof. For shortness, by a denote the point $a_1 = a_2$, and by P the piece of the boundary of R from b_1 to b_2 according to the positive orientation, so this piece which does not contain a.

Let $x \in P$ be an extreme point of R different from b_1 and b_2. By Theorem 4 of [12] there exists a lune L of thickness $\Delta(R)$ containing R such that x is the center of one of the semicircles bounding L. The lune L has the form $K \cap K^*$, where K^* is the (unique, by Part I of Theorem 1 of [12]) hemisphere supporting R at x. Since x lays strictly between b_1 and b_2 on the boundary of R and since the different points b_1, x, b_2 are the centers of the semicircles $M_1^*/M_1, K^*/K, M_2^*/M_2$, respectively, we conclude that the hemisphere K^* satisfies $\prec M_1^* K^* M_2^*$. Thus, by Lemma 2 we obtain $\prec M_1 K M_2$ and, as a consequence, a is the only point of support of R by K. By Claim 2 of [12] the center of K/K^* belongs to R, which implies that this center is at the point a. Therefore $|ax| = \Delta(R)$. Since x is an arbitrary extreme point of P, we conclude that every extreme point of P is in the distance $\Delta(R)$ from a.

Assume for a while that P contains an arc x_1x_2. By the preceding paragraph, x_1 and x_2 are in the distance $\Delta(R)$ from a. Observe that all points laying strictly inside this arc are in the distance less than $\Delta(R)$ from a. Let $H(o)$ be the hemisphere supporting R such that the arc oa intersects the arc x_1x_2 in the point y different from x_1 and x_2. Since $|ay| < \Delta(R)$, the disk centered at o with radius $\frac{\pi}{2} - \Delta(R)$ does not touch R. Therefore by Theorem 1 of [12] the lune $H(o) \cap H(o)^*$ is of thickness less than $\Delta(R)$. It contradicts the fact that R has thickness $\Delta(R)$. Thus we conclude that P does not contain any arc.

From the two preceding paragraphs we conclude that all points of b_1b_2 are extreme and all of them are in the distance $\Delta(R)$ from a. Consequently, we obtain the first thesis of our proposition

For the proof of the second part of this proposition take any hemisphere M satisfying $\prec M_1 M M_2$. Denote by B the disk of radius $\frac{\pi}{2} - \Delta(R)$ concentric with M. By Part I of Theorem 1 from [12] there is a unique hemisphere M^* and the point of supporting R by M^* is the point where B touches R. Moreover, $\Delta(M \cap M^*) = \frac{\pi}{2} - \left(\frac{\pi}{2} - \Delta(R)\right) = \Delta(R)$. So we get the second part of our proposition. □

Theorem 2. *Let $R \subset S^2$ be a reduced body with $\Delta(R) < \frac{\pi}{2}$. Assume that M is a supporting hemisphere of R such that the intersection of* $\mathrm{bd}(M)$ *with* $\mathrm{bd}(R)$ *is a non-degenerated arc x_1x_2. Then* $\mathrm{width}_M(R) = \Delta(R)$*, and the center of M/M^* belongs to x_1x_2.*

Proof. First, let us show that $\mathrm{width}_M(R) = \Delta(R)$. Assume the opposite, i.e., that $\mathrm{width}_M(R) > \Delta(R)$. Then applying the continuity of the width (see Theorem 2 of [12]), we conclude that there exist supporting hemispheres M_1 and M_2 of R such that $\mathrm{width}_{M_1}(R) = \Delta(R) = \mathrm{width}_{M_2}(R)$ and $\mathrm{width}_M(R) > \Delta(R)$ for every hemisphere M supporting R strictly between M_1 and M_2. Clearly, M_1 supports R at x_1 and M_2 supports R at x_2. Denote by x_0 the intersection point of the semicircles M_1/M_1^* and M_2/M_2^*. Since $x_0 \notin x_1x_2$ and $x_1x_2 \subset \mathrm{bd}(R)$, we have $x_0 \notin R$. But on the other hand, by the result of the paragraph before the last of the proof of Theorem 1 we know that $x_1 = x_0$ or $x_2 = x_0$, which means that $x_0 \in R$. This contradiction implies that $\mathrm{width}_M(R) = \Delta(R)$.

The second part of our theorem is obvious by $(M/M^*) \cap R = x_1x_2$ and since from Claim 2 in [12] we know that the center of M/M^* belongs to R. □

Theorem 3. *If M is an extreme supporting hemisphere of a reduced spherical body $R \subset S^2$, then* $\mathrm{width}_M(R) = \Delta(R)$.

Proof. Let M be an extreme supporting hemisphere of R, say right, at a point p. If the great circle bounding M contains more than one point of $\mathrm{bd}(R)$, we apply Theorem 2. Otherwise there is a sequence $\{p_i\}$ of different points of $\mathrm{bd}(R)$ monotonically convergent to p from the right and such that all the arcs pp_i are not in $\mathrm{bd}(R)$. The great circle containing p and p_i dissects R into two closed convex subsets. Denote by R_i this of these subsets which does not contain p_{i+1} and by M_i the hemisphere supporting R_i at the points of the arc pp_i. Since R is reduced, $\mathrm{width}_M(R_i) < \Delta(R)$. The lune $M_i \cap M_i^*$ has thickness $\mathrm{width}_M(R_i)$ and the limit of

lunes $M_i \cap M_i^*$, as i tends to ∞, is the lune $M \cap M^*$. Hence the thickness of $M \cap M^*$, which is equal to $\mathrm{width}_M(R)$, cannot be greater than $\Delta(R)$. Consequently, it is equal to $\Delta(R)$. □

Proposition 2. *Let p be a point of a reduced body $R \subset S^2$ of thickness at most $\frac{\pi}{2}$. Then $R \subset H(p)$.*

Proof. If p is an extreme point of R, then by the two-dimensional case of Theorem 4 from [12] there exists a lune $L \supset R$ of thickness $\Delta(R)$ with p as the center of one of the semicircles bounding L. Observe that $L \subset H(p)$. Hence $R \subset H(p)$, which ends the proof in this case.

If p is a boundary but not an extreme point of $\mathrm{bd}(R)$, then there exist extreme points e_1, e_2 of R such that $p \in e_1e_2$. We already know that $R \subset H(e_i)$ for $i = 1, 2$. Thus $R \subset H(e_1) \cap H(e_2)$. Observe that both the semicircles bounding the lune $H(e_1) \cap H(e_2)$ are contained in $H(p)$. Hence this lune is contained in $H(p)$. Therefore $R \subset H(p)$, which ends the proof in this case.

If p is not a boundary point of R, then there exist boundary points b_1, b_2 of R such that $p \in b_1b_2$. For analogical reason as in the preceding paragraph, we have $R \subset H(b_1) \cap H(b_2) \subset H(p)$, which ends the proof. □

This Proposition is applied in the proofs of Theorems 4 and 7.

4 Reduced bodies of thickness at least $\frac{\pi}{2}$

Proposition 3 and Theorem 4 say on the shape of reduced bodies of thickness at least $\frac{\pi}{2}$. The proof of first of them requires the following lemma.

Lemma 3. *Let $C \subset S^d$ be a spherical convex body with $\Delta(C) > \frac{\pi}{2}$ and let $L \supset C$ be a lune such that $\Delta(L) = \Delta(C)$. Each of the centers of the $(d-1)$-dimensional hemispheres bounding L belongs to the boundary of C and both they are smooth points of the boundary of C.*

Proof. We have $L = K \cap M$, where K and M are two hemispheres. Both of them must support C, because in the opposite case we could find a narrower lune than L containing C, which, by the assumption $\Delta(L) = \Delta(C)$, is impossible. For instance, let us show the thesis of our lemma for the $(d-1)$-dimensional hemisphere M/K bounding L. Since $\Delta(C) > \frac{\pi}{2}$, according to Part III of Theorem 1 from [12] the hemisphere M is a (possibly non-unique) hemisphere fulfilling $M = K^*$. Denote by k the center of K. After the just mentioned Part III, the hemisphere M supports C at exactly one point t of the set $\mathrm{bd}(C) \cap B$, where B denotes the largest ball with center k contained in C. By Corollary 2 of [12] the point t is the center of M/K. Since t belongs to $\mathrm{bd}(C) \cap B$, it is a smooth point of $\mathrm{bd}(C)$, which ends the proof. □

Proposition 3. *Every reduced spherical body $R \subset S^d$ of thickness over $\frac{\pi}{2}$ is smooth.*

Proof. In order to see this recall that for every extreme point e of R there exists a lune L as in Theorem 4 of [12] fulfilling $\Delta(L) = \Delta(R)$, so that e is the center of one of the semicircles bounding L. By Lemma 3 we obtain that e is a smooth point of the boundary of R. Since every extreme point of R is a smooth boundary point of R, we conclude that R is a smooth spherical convex body. □

The thesis of Proposition 3 does not hold true for every thickness at most $\frac{\pi}{2}$. For instance, on S^2 the regular spherical triangle of any thickness at most $\frac{\pi}{2}$ is reduced but not smooth.

Theorem 4. *Every reduced spherical convex body $R \subset S^2$ such that $\Delta(R) \geq \frac{\pi}{2}$ is a spherical body of constant width $\Delta(R)$.*

Proof. If $\Delta(R) > \frac{\pi}{2}$, we apply Proposition 3. By Theorem 5 of [12], which says that every smooth reduced spherical body is a body of constant width, we obtain the thesis of our theorem.

Consider the case when $\Delta(R) = \frac{\pi}{2}$. Assume that there exists a hemisphere K supporting R such that for a fixed K^* the lune $L = K \cap K^*$ has the thickness over $\frac{\pi}{2}$. From Corollary 2 in [12] we obtain that the center of the semicircle K^*/K belongs to R. Denote it by b.

Clearly, there exists an extreme point e of R on the semicircle K/K^*. By Proposition 2 the body R is contained in the hemisphere M with the center at e. Thus $|eb| \leq \frac{\pi}{2}$. But since $\Delta(L) > \frac{\pi}{2}$, the only points of K/K^* lying in the distance at most $\frac{\pi}{2}$ from b are the corners of L. It means that e is a corner of L and moreover it is the only point of $K/K^* \cap R$.

Observe that since e lies on the boundary of K, the lune $K \cap M$ has the thickness $\frac{\pi}{2}$. From the definition of $\mathrm{width}_K(R)$ we obtain that the lune $K \cap K^*$ is of thickness at most $\Delta(K \cap M) = \frac{\pi}{2}$ which contradicts our assumption. Thus R is of constant width. □

5 The case of spherical bodies of constant width

Recall that every spherical convex body of constant width is reduced. Theorem 4 says that the opposite implication holds true for bodies of thickness at least $\frac{\pi}{2}$ on S^2, i.e. every reduced spherical convex body of thickness at least $\frac{\pi}{2}$ on S^2 is of constant width. Observe that it does not hold true for reduced spherical convex bodies of thickness below $\frac{\pi}{2}$. For example every regular spherical triangle of thickness below $\frac{\pi}{2}$ is reduced but is not of constant width.

Theorem 5. *Every spherical strictly convex reduced body of thickness smaller than $\frac{\pi}{2}$ on S^2 is of constant width.*

Proof. Assume that a spherical strictly convex body R of thickness less than $\frac{\pi}{2}$ is reduced but not of constant width. It means that there exists a hemisphere M_0 for which $\mathrm{width}_{M_0}(R) > \Delta(R)$. Due to continuity arguments, there exist hemispheres M_1 and M_2 supporting R such that $\mathrm{width}_{M_1}(R) = \Delta(R) = \mathrm{width}_{M_2}(R)$ and $\mathrm{width}_M(R) > \Delta(R)$ for every hemisphere M supporting R strictly between M_1 and M_2. From Theorem 1 we see that the arc with the end-points at the centers of the semicircles M_1/M_1^* and M_2/M_2^* belongs to the boundary of R. Thus R is not strictly convex. □

Here is, in a sense, an opposite theorem.

Theorem 6. *Every spherical convex body of constant width smaller than $\frac{\pi}{2}$ on S^2 is strictly convex.*

Proof. Assume the opposite, i.e., that there exists a body W of constant width below $\frac{\pi}{2}$ and different extreme points e_1, e_2 of W such that the arc e_1e_2 belongs to $\mathrm{bd}(W)$.

Denote by N_1 the hemisphere supporting W at every point of e_1e_2. Clearly, its bounding great circle contains e_1e_2. Since W is of constant width, the lune $L_1 = N_1 \cap N_1^*$ has thickness $\Delta(W)$, and since W, as a body of constant width, is a reduced body, by Theorem 2 the center c of the semicircle N_1/N_1^* belongs to e_1e_2. Since c must be different from e_1 or e_2, for instance assume that c is not at e_1.

In the preceding and the next paragraph the hemispheres N_1^* and N_2^* supporting W are unique by Part I of Theorem 1 from [12]. The reason is that W is of constant width below $\frac{\pi}{2}$.

By Theorem 4 of [12] there exists a hemisphere N_2 supporting W at e_1 such that e_1 is the center of N_2/N_2^*. Assume for a while that $N_1 = N_2$. From the preceding paragraph we conclude that $N_1^* = N_2^*$ and so $N_1 \cap N_1^* = N_2 \cap N_2^*$. Therefore the center c of N_1/N_1^* coincides with the center e_1 of N_2/N_2^*. It contradicts $c \neq e_1$ and thus we obtain that the hemispheres N_1 and N_2 must be different.

Denote by o_i the center of N_i for $i = 1, 2$. Observe that every hemisphere supporting W strictly between N_1 and N_2, supports W only at e_1. Moreover, the center of every such a hemisphere lays on the arc o_1o_2 and vice-versa, every point of this arc is the center of a hemisphere supporting W strictly between N_1 and N_2.

Let t, u be points in the distance $\Delta(W)$ from e_1 such that $t \in e_1o_1$ and $u \in e_1o_2$. By Corollary 2 of [12] we see that u is the center of N_2^*/N_2 and it belongs to W. Observe that t belongs to the disk of radius $\frac{\pi}{2} - \Delta(W)$ with the center at o_1. By Part I of Theorem 1 of [12] this disk touches W only at the center of N_1^*/N_1. Denote this center by c^*. Since the point c^* lays on the arc co_1, the point t lays on the arc e_1o_1 and the only common point of co_1 and e_1o_1 is o_1, we see that c^* is different from t. Therefore t does not belong to W.

Consider the shorter part P of the spherical circle with the center at e_1 and radius $\Delta(W)$ between t and u. Since W is closed, $t \notin W$ and $u \in W$, we conclude that there exists a point $x \in P$ different from t and not contained in W. Denote by o the point of the arc o_1o_2 such that $x \in oe_1$ and by N the hemisphere centered at o. Since o lays strictly between o_1 and o_2, the hemisphere N supports W strictly between N_1 and N_2. As a consequence, N supports W only at e_1. Thus by Claim 2 of [12] the center of N/N^* is at e_1.

Looking to the proof of Part I of Theorem 1 from [12], we observe that the center of our N^*/N lays on the arc connecting the center of N with the center of N/N^* in the distance $\Delta(W)$ from the center of N/N^*. Therefore x is the center of N^*/N and by Claim 2 of [12] it belongs to the boundary of W. But this contradicts the fact that x does not belong to W. This shows that the assumption from the beginning of our proof must be false. Hence our theorem is proved. □

The thesis of this Theorem does not hold true for spherical convex bodies of constant width at least $\frac{\pi}{2}$, as we see from the following Example.

Example. Take a spherical regular triangle abc of sides of length $\kappa < \frac{\pi}{2}$ and prolong them by the same distance $\sigma \leq \frac{\pi}{2} - \kappa$ in both "directions" up to points d, e, f, g, h, i, so that i, a, b, f are on a great circle in this order, that e, b, c, h are on a great circle in this order, and that g, c, a, d are on a great circle in this order. Provide three pieces of circles of radius $\kappa + \sigma$: with center a from f to g, with center b from h to i, with center c from d to e. Moreover three pieces of circles of radius σ: with center a from i to d, with center b from e to f, with center c from g to h. The convex hull U of these six pieces of circles is a spherical body of constant width $\kappa + 2\sigma$. In particular, when $\kappa + \sigma = \frac{\pi}{2}$, three boundary circles of U become arcs; namely de, fg and hi.

In [12] the question is asked if through every boundary point p of a reduced spherical body R a lune $L \supset R$ of thickness $\Delta(R)$ passes with p as the center of one of the two semicircles bounding L. The example of the regular spherical triangle of thickness less than $\frac{\pi}{2}$ shows that the answer is negative in general. The following proposition shows that for bodies of constant width the answer is positive.

Theorem 7. *Let $W \subset S^2$ be a spherical body of constant width. For every boundary point p of W there exists a lune $L \supset W$ fulfilling $\Delta(L) = \Delta(W)$ such that p is the center of one of the semicircles bounding L.*

Proof. If p is an extreme point of W, then for this point the thesis is true by Theorem 4 of [12]. In particular, the thesis is true for bodies of constant width below $\frac{\pi}{2}$, since then by Theorem 6 all boundary points of W are extreme.

Consider the case when $\Delta(W) = \frac{\pi}{2}$ and p is not an extreme point of W. Clearly, there is a unique hemisphere G supporting W at p. By Proposition 2 we have $W \subset H(p)$. Consider the lune $H(p) \cap G$ and note that it contains W. Observe that p is in the distance $\frac{\pi}{2}$ from its



FIGURE 2. Illustration to the proof of Theorem 7.

corners, and thus it is the center of a semicircle bounding it. Moreover it is in the distance $\frac{\pi}{2}$ from every point of the other semicircle bounding this lune. Hence this lune is of thickness $\Delta(W)$. Thus $H(p) \cap G$ is a lune promised in our theorem.

Finally take into account the case when $\Delta(W) > \frac{\pi}{2}$ and p is not an extreme point of W (the just presented Example shows that it may happen), and let us stay with this assumption up to the end of the proof. Clearly, p belongs to a boundary arc e_1e_2 of the body W, where e_1 and e_2 are extreme points of W (see Figure 2).

Denote by K the hemisphere supporting W whose bounding great circle contains the arc e_1e_2 and by k denote the center of K. From Proposition 3 we know that K is the only hemisphere supporting W at each of the points e_1 and e_2. By Theorem 4 of [12] there exist unique hemispheres K_1^* and K_2^* (they play the part of K^* in Theorem 1 of [12]) supporting W such that the lunes $K \cap K_1^*$ and $K \cap K_2^*$ are of thickness $\Delta(W)$ with e_1 as the center of K/K_1^*, and e_2 as the center of K/K_2^* (e_i plays the part of s in Part III of Theorem 1 from [12]). Denote by t_1 the center of K_1^*/K and by t_2 the center of K_2^*/K (so t_i plays the part of t in the proof of Part III of Theorem 1 from [12]). Clearly the disk B of radius $\rho = \Delta(W) - \frac{\pi}{2}$ and with the center at k (as in Part III of Theorem 1 in [12]) touches W from inside at the points t_1 and t_2. Moreover, from the proof of Theorem 1 of [12] and from the earlier established fact that e_i is the center of K/K_i^* and t_i the center of K_i^*/K, where $i = 1, 2$, we obtain that k belongs to both arcs e_1t_1 and e_2t_2.

Assume for a while that the shorter piece P of $\mathrm{bd}(B)$ between t_1 and t_2 is not in $\mathrm{bd}(W)$. Take the corresponding piece of $\mathrm{bd}(W)$ between t_1 and t_2. Clearly, there is an extreme point $x \notin P$ of this piece. The great circle containing xk intersects P at a point y, and thus also e_1e_2 at a point q. Clearly, $|xq| > |yq| = |yk| + |kq| = \rho + \frac{\pi}{2} = \Delta(W)$. By Theorem 4 of [12] there exist hemispheres G and H such that the lune $J = G \cap H$ of thickness $\Delta(W)$ contains W with x as the center of H/G. Denote by z the center of G/H. Since $\Delta(J) > \frac{\pi}{2}$, the only farthest point from x on G is just z. Hence $|xq| \leq |xz| = \Delta(J) = \Delta(W)$. A contradiction with $|xq| > \Delta(W)$ established earlier. Consequently, $P \subset \mathrm{bd}(W)$.

Observe that the great circle through p and k intersects P at a point r different from t_1 and t_2. Since k is the center of the hemisphere K, we see that kp is orthogonal to the

semicircle K/N. Since k is also the center of the disk B and since r is the point at which the great circle N touches B, we see that kr is orthogonal to the semicircle N/K. From these two observations we obtain that pr is orthogonal to both the semicircles K/N and N/K. Thus p and r are the centers of the semicircles bounding L (it follows from the obvious fact that the only orthogonal great circle to two different non-opposite meridians is the equator). Moreover, L is of thickness $|pr| = \Delta(W)$ and hence L is the lune announced in our theorem. □

6 Diameter of reduced spherical bodies

In the next theorem we prove the conjecture presented in the paragraph before the last of [13]. For the proof we need the following two lemmas. By the way, both may be easily generalized for S^d, but for our needs the case when $d = 2$ is sufficient.

Lemma 4. *Let $L \subset S^2$ be a lune of thickness at most $\frac{\pi}{2}$ whose bounding semicircles are Q and Q'. For every u, v, z in Q such that $v \in uz$ and for every $q \in L$ we have $|qv| \leq \max\{|qu|, |qz|\}$.*

Proof. If $\Delta(L) = \frac{\pi}{2}$ and q is the center of Q', then the distance between q and any point of Q is the same, and thus the thesis is obvious. Consider the opposite case, i.e., when $\Delta(L) < \frac{\pi}{2}$, or $\Delta(L) = \frac{\pi}{2}$ but q is not the center of Q'. Clearly, the closest point $p \in Q$ to q is unique. Observe that for $x \in Q$ the distance $|qx|$ increases as the distance $|px|$ increases. This easily implies the thesis of our lemma. □

For a convex body $C \subset S^2$ denote by $\operatorname{diam}(C)$ its diameter and by $E(C)$ the set of its extreme points.

Lemma 5. *For every spherical convex body $C \subset S^2$ of diameter at most $\frac{\pi}{2}$ we have* $\operatorname{diam}(E(C)) = \operatorname{diam}(C)$.

Proof. Clearly, $\operatorname{diam}(E(C)) \leq \operatorname{diam}(C)$. In order to show the opposite inequality $\operatorname{diam}(C) \leq \operatorname{diam}(E(C))$, thanks to $\operatorname{diam}(C) = \operatorname{diam}(\operatorname{bd}(C))$, it is sufficient to show that $|cd| \leq \operatorname{diam}(E(C))$ for any $c, d \in \operatorname{bd}(C)$. If $c, d \in E(C)$, this is trivial. In the opposite case, at least one of these points does not belong to $E(C)$. If, say $d \notin E(C)$, then by $d \in \operatorname{bd}(C)$ there are $e, f \in E(C)$ different from d such that $d \in ef$. This and $d, e, f \in \operatorname{bd}(C)$ imply that the arc ef is a subset of $\operatorname{bd}(C)$.

Recall that by Theorem 3 from [13] we have $\operatorname{width}_K(C) \leq \operatorname{diam}(C)$ for every hemisphere K supporting C. In particular, for the hemisphere supporting C at every point of the arc ef. Thus by the assumption that $\operatorname{diam}(C) \leq \frac{\pi}{2}$ we obtain $\operatorname{width}_K(C) \leq \frac{\pi}{2}$. Hence we may apply Lemma 4. We obtain $|cd| \leq \max\{|ce|, |cf|\}$.

If $c \in E(C)$, from $e, f \in E(C)$ we conclude that $|cd| \leq \operatorname{diam}(E(C))$. If $c \notin E(C)$, from $c \in \operatorname{bd}(C)$ we see that there are $g, h \in E(C)$ such that $c \in gh$. Similarly as in the consideration of the preceding paragraph we see that $|ec| \leq \max\{|eg|, |eh|\}$ and $|fc| \leq \max\{|fg|, |fh|\}$. And again by the inequality at the end of the preceding paragraph and by these two just shown inequalities we get $|cd| \leq \max\{|eg|, |eh|, |fg|, |fh|\} \leq \operatorname{diam}(E(C))$, which ends the proof. □

The assumption that $\operatorname{diam}(C) \leq \frac{\pi}{2}$ is substantial in this lemma, as it follows from the example of any regular triangle of diameter over $\frac{\pi}{2}$. The weaker assumption that $\Delta(C) \leq \frac{\pi}{2}$ is not sufficient, which follows from the example of any isosceles triangle T with $\Delta(T) \leq \frac{\pi}{2}$ and the arms longer than $\frac{\pi}{2}$ (so the base shorter than $\frac{\pi}{2}$). The diameter of T equals to the distance between the midpoint of the base and the opposite vertex of T. Hence $\operatorname{diam}(T)$ is over the length of each of the sides.

Theorem 8. *For every reduced spherical body $R \subset S^2$ with $\Delta(R) < \frac{\pi}{2}$ we have* $\operatorname{diam}(R) \leq \arccos(\cos^2 \Delta(R))$. *This value is attained if and only if R is the quarter of disk of radius $\Delta(R)$. If $\Delta(R) \geq \frac{\pi}{2}$, then* $\operatorname{diam}(R) = \Delta(R)$.



FIGURE 3. Illustration to the proof of Theorem 8.

Proof. Assume that $\Delta(R) < \frac{\pi}{2}$. By Lemma 5 it is sufficient to show that the distance between any two points of $E(R)$ is at most $\arccos(\cos^2 \Delta(R))$. Let e_1, e_2 be different extreme points of R. Since R is reduced, according to the statement of Theorem 4 from [12] there exist lunes $L_j \supset R$, where $j = 1, 2$, of thickness $\Delta(R)$ with e_j as the center of one of two semicircles bounding L_j (see Figure 3). Denote by b_j, where $j = 1, 2$, the center of the other semicircle bounding L_j.

If the lunes L_1 and L_2 coincide and $e_1 = b_2$, $e_2 = b_1$, then $|e_1e_2| = \Delta(R) \leq \arccos(\cos^2 \Delta(R))$. Otherwise $L_1 \cap L_2$ is a spherical quadrangle with points e_1, b_2, b_1, e_2 on the consecutive sides. Therefore e_1b_1 and e_2b_2 intersect at exactly one point. Denote it by g. Provide the great circle F orthogonal to e_1b_1 which passes through e_2. Since $e_2 \in L_1$, we see that F intersects e_1b_1. Let f be the intersection point of them. From $|e_2b_2| = \Delta(R)$ we see that $|ge_2| \leq \Delta(R)$. Thus from the right triangle gfe_2 we conclude that $|fe_2| \leq \Delta(R)$. Moreover, from $|e_1b_1| = \Delta(R)$ and $f \in e_1b_1$ we see that $|fe_1| \leq \Delta(R)$. Consequently, from the formula $\cos k = \cos l_1 \cos l_2$ for a right spherical triangle with hypotenuse k and legs l_1, l_2 applied to the triangle e_1fe_2 (again see Figure 3) we obtain $|e_1e_2| \leq \arccos(\cos^2 \Delta(R))$.

Observe that if $\Delta(R) < \frac{\pi}{2}$, then the shown inequality becomes the equality only if $g = f = b_1$ and $|b_1e_2| = \Delta(R)$. In this case, by Proposition 1, our body R is a quarter of disk of radius $\Delta(R)$.

Finally, when we intend to show the last statement of our theorem, assume that $\Delta(R) \geq \frac{\pi}{2}$. By Theorem 4, the body R is of constant width $\Delta(R)$.

Take a diametral segment pq of R. Clearly, $p \in \mathrm{bd}(R)$. Take the lune L from Theorem 7 such that p is the center of a semicircle bounding L. Denote by s the center of the other semicircle S bounding L. By the third part of Lemma 3 of [12], we have $|px| < |ps|$ for every $x \neq s$ on S. Hence $|px| \leq |ps|$ for every $x \in S$. So also $|pz| \leq |ps|$ for every $z \in L$. In particular, $|pq| \leq |ps|$. Since $\mathrm{diam}(R) = |pq|$ and $\Delta(R) = \Delta(L) = |ps|$, we obtain $\mathrm{diam}(R) \leq \Delta(R)$.

On the other hand, $\Delta(R) \leq \mathrm{diam}(R)$ by Proposition 1 of [12]. Hence the last statement of our theorem is true. □

Let us add that an analogical theorem on the diameter of spherical reduced polygons is presented in [13].

Now, in this section on the diameter, let us repeat Proposition 2 in the following form: *for every reduced spherical body with* $\Delta(R) \leq \frac{\pi}{2}$ *on* S^2 *we have* $\mathrm{diam}(R) \leq \frac{\pi}{2}$. Here is even a more precise form of it.

Proposition 4. *Let $R \subset S^2$ be a reduced spherical body. We have $\Delta(R) < \frac{\pi}{2}$ if and only if* $\operatorname{diam}(R) < \frac{\pi}{2}$. *Moreover, $\Delta(R) = \frac{\pi}{2}$ if and only if* $\operatorname{diam}(R) = \frac{\pi}{2}$.

Proof. Applying the first derivative test, we see that $f(x) = \arccos(\cos^2 x)$ is an increasing function in the interval $\left[0, \frac{\pi}{2}\right]$. Moreover, for $x = \frac{\pi}{2}$ this function accepts the value $\frac{\pi}{2}$. So in the interval $[0, \frac{\pi}{2})$ it accepts only the values below $\frac{\pi}{2}$. Thus by Theorem 8, if $\Delta(R) < \frac{\pi}{2}$, then $\operatorname{diam}(R) < \frac{\pi}{2}$. From the fact that $\Delta(C) \leq \operatorname{diam}(C)$ for any spherical convex body C, which follows from Theorem 3 and Proposition 1 of [12], we obtain the opposite implication. The second statement is an easy consequence of Proposition 2 and the first statement of our proposition. □

References

[1] L. Danzer, B. Grünbaum and V. Klee, Helly's theorem and its relatives, in Proc. of Symp. in Pure Math. vol. VII, Convexity, 1963, pp. 99–180.

[2] L. Fejes Tóth, *Lagerungen in der Ebene auf der Kugel und im Raum.* (German) Zweite verbesserte und erweiterte Auflage. Die Grundlehren der mathematischen Wissenschaften, Band 65. Springer-Verlag (Berlin-New York, 1972).

[3] L. Fejes Tóth Toth, Research Problems: Exploring a Planet. *Amer. Math. Monthly* **80** (1973), 1043-1044.

[4] O. P. Ferreira, A. N. Iusem and S. Z. Németh, Projections onto convex sets on the sphere, J. Global Optim. **57** (2013), 663–676.

[5] F. Gao, D. Hug and R. Schneider, Intrinsic volumes and polar sets in spherical space, Homage to Luis Santaló. Vol. 1 Math. Notae **41** (2001/02), 159–176.

[6] H. Groemer, Extremal convex sets, *Monatsh. Math.* **96** (1983), 29–39.

[7] H. Hadwiger, Kleine Studie zur kombinatorischen Geometrie der Sphäre, *Nagoya Math. J.* **8** (1955), 45–48.

[8] H. Han and T. Nishimura, Self-dual shapes and spherical convex bodies of constant width $\pi/2$, *J. Math. Soc. Japan*, to appear (arXiv:1511.04165v2).

[9] E. Heil, *Kleinste konvexe Körper gegebener Dicke.* Preprint No. 453, Fachbereich Mathematik der TH Darmstadt, 1978.

[10] E. Heil and H. Martini, Special convex bodies, in Handbook of Convex Geometry, *North-Holland, Elsevier Science Publishers*, pp. 347–385 (1993)

[11] M. Lassak, Reduced convex bodies in the plane, *Israel J. Math.* **70** (1990), 365–379.

[12] M. Lassak, Width of spherical convex bodies, *Aequationes Math.* **89** (2015), 555–567.

[13] M. Lassak, Reduced spherical polygons, *Colloq. Math.* **138**(2015), 205–216.

[14] M. Lassak and H. Martini, Reduced convex bodies in Euclidean space – a survey, *Expositiones Math.* **29** (2011), 204–219.

[15] Y. S. Lee, J. M. Yang and I. K. Park, Convex figures on the unit sphere, *Far East J. Math. Sci.* **7** (2002), 191–196

[16] K. Leichtweiss, Curves of constant width in the non-Euclidean geometry, *Abh. Math. Sem. Univ. Hamburg* **75** (2005), 257–284.

[17] T. Nishimura, Y. Sakemi, Topological aspects of Wulff shapes, *J. Math. Soc. Japan* **66** (2014) 89-109.

[18] G. Van Brummelen, *Heavenly mathematics. The forgotten art of spherical trigonometry.* Princeton University Press (Princeton, 2013)